

RESEARCH

Open Access



Survey of farm, parlour and milking management, parlour technologies, SCC control strategies and farmer demographics on Irish dairy farms

Alice Uí Chearbhaill^{1,2*} , Pablo Silva Boloña¹, Eoin G. Ryan², Catherine I. McAloon², Alison Burrell³, Conor G. McAloon² and John Upton¹

Abstract

Background This cross-sectional study describes a survey designed to fill knowledge gaps regarding farm management practices, parlour management practices and implemented technologies, milking management practices, somatic cell count (SCC) control strategies, farmer demographics and attitudes around SCC management on a sample of Irish dairy farms.

Results We categorized 376 complete responses by herd size quartile and calving pattern. The average respondent herd was 131 cows with most (82.2%) operating a seasonal calving system. The median monthly bulk tank somatic cell count for seasonal calving systems was 137,000 cells/ml (range 20,000 – 1,269,000 cells/ml), 170,000 cells/ml for split-calving systems (range 46,000 – 644,000 cells/ml) and 186,000 cells/ml for ‘other’ herds (range 20,000 – 664,000 cells/ml). The most common parlour types were swing-over herringbones (59.1%) and herringbones with recording jars (22.2%). The average number of units across herringbone parlours was 15, 49 in rotary parlours and two boxes on automatic milking system (AMS) farms. The most common parlour technologies were in-parlour feeding systems (84.5%), automatic washers on the bulk tank (72.8%), automatic cluster removers (57.9%), and entrance or exit gates controlled from the parlour pit (52.2%). Veterinary professionals, farming colleagues and processor milk quality advisors were the most commonly utilised sources of advice for SCC management (by 76.9%, 50.0% and 39.2% of respondents respectively).

Conclusions In this study, we successfully utilised a national survey to quantify farm management practices, parlour management practices and technology adoption levels, milking management practices, SCC control strategies and farmer demographics on 376 dairy farms in the Republic of Ireland. Rotary and AMS parlours had the most parlour technologies of any parlour type. Technology add-ons were generally less prevalent on farms with smaller herds. Despite finding areas for improvement with regard to frequency of liner changes, glove-wearing practices and engagement with bacteriology of milk samples, we also found evidence of high levels of documentation of mastitis treatments and high use of post-milking teat disinfection. We discovered that Irish dairy farmers are relatively content in their careers but face pressures regarding changes to the legislation around prudent antimicrobial use in their herds.




Fig. 2 Median monthly BTSCC by herd size quartiles across 12 months of 2021 and 8 months of 2022 (from processor dataset) ($n=376$)




Fig. 3 Median monthly BTSCC by calving patterns across 12 months of 2021 and 8 months of 2022 (from processor dataset) ($n=376$). ‘SEASONAL’ refers to herds which calved all cows between February and April and those in which peak milk production occurred in May or June and exceeded the herd’s minimum monthly milk production in the herd year by $>700\%$ ($n=309$). ‘SPLIT’ refers to herds which supplied milk throughout December and January and had peak milk production that exceeded herd minimum milk production for any month by $<300\%$ ($n=29$). ‘OTHER’ calving patterns are milk producers which do not fit into either of the two former categories due to the limitations set in characterising them ($n=38$)

Farm technology and management survey

Farm management

Herds managed as part of a partnership accounted for 22.3% of herds and 3.7% of respondents milked in more than one parlour over the course of lactation.

Most respondents milked twice per day (95.2%), 1.3% of respondents milked once per day, and 0.8% responded with ‘other’. The ‘other’ answers pertained to different milking practices at different times of the year, for example “13 milkings [from] July to Dec” and “do 10 in 7 [days] for




Fig. 4 Percentage of respondent herds with monthly BTSCC equal to or below 100,000 cells/ml, between 101–200,000 cells/ml, between 201–399 cells/ml and equal to or above 400,000 cells/ml across 12 months of 2021 and 8 months of 2022 (from processor dataset) ($n = 376$)

the last three months of 2021". Respondents that milked with an AMS and therefore had no set milking schedule accounted for 2.7% of responses.

Morning milking most commonly started between 7 and 8 am (61.0% of respondents) and most commonly lasted 90 min (25.2%). Evening milking start time most commonly occurred between 4 and 5 pm (55.0%) and lasted 90 min (30.0%). The most common interval between morning and evening milkings was 9 to 10 h (60.1%). Most commonly, there was only one person milking in the parlour for morning (68.8%) and evening (66.5%) milkings. During milking, 90.7% of the survey respondents were present at the milking themselves, 57.1% enlisted a family member and 33.6% hired an employee. Most of the milking parlours were less than 10 years of age (46.8%), with 20.1% of these being less than 5 years of age. Upgrades were applied to parlours in the past five years on 36.8% of respondent farms.

Farmers were asked for their most likely influencing factors for culling on a 7-point scale ('1' being most likely reason to cull and '7' being least likely reason to cull). Fertility problems were given a score of '1' or '2' by 50.2% of respondent farms, as was persistent high SCC (45.0%) and recurrent clinical mastitis (40.7%). Lameness and poor milk production were awarded mid-level scores of '3–5' by 60.8% and 57.6% of respondent farms respectively. A score of '6' or '7' was awarded to

individual cow behaviour by 56.0% and age by 53.4% of respondent farms.

Parlour management and technologies

The most prevalent parlour types were swing-over herringbones (59.1%) and herringbones with milk recording jars (22.2%). This was followed by parallel parlours (6.7%), double-up herringbones (4.8%), rotaries (2.7%) and AMS (2.7%). Less frequent parlour types submitted as an 'other' option included one abreast parlour, one bucket plant, two herringbones with no recording jars, one parallel parlour with recording jars, one non-descript herringbone and a rapid exit parlour. The average number of units across the various types of herringbone parlours was 15 (range 2–40), 49 in rotary parlours (range 40–70) and two boxes on AMS farms (range 1–3). Most parlours had a mid-level milk line (90.1%). The most common pulsation system was an alternating system (53.9%) followed by simultaneous (41.3%) and 'one pulsator per teat' systems (3.8%).

One parlour service per year was carried out on 72.1% of respondent farms, 15.2% serviced twice per year, 8.0% serviced less than once per year and 4.8% serviced more than twice per year. Liner changes occurred most commonly once (34.4%) or twice (44.8%) per year. Other variations of liner change frequency included every 2,000

milking (6.7%), every 2,500 milking (6.1%), three times per year (4.5%) or every two years (1.3%).

Farmers had various means of ensuring good cow positioning in the parlour. A full list of cow positioning additions by parlour type can be found in Supplementary Table 6 and a list of total positioning additions by parlour type in Supplementary Table 7. Individual mangers were present in 59.6% of parlours. Straight rump rails were present in 39.3% of parlours whilst 14.8% had zig-zag rump rails. Manual bailing systems were present in 13.7% of respondent parlours whilst 5.5% had a sequential bailing system. Straight breast rails and adjustable breast rails were present in 7.1% and 8.5%, respectively, of respondent parlours.

Farmers also had varying degrees of technological additions in their parlours. A full list can be observed in Fig. 5. A full list of parlour technology additions by herd size quartiles can be found in Table 3 and Supplementary Table 4. A full list of parlour technology additions by parlour type can be observed in Supplementary Table 8 and a list of total technology additions by parlour type in Supplementary Table 5. No technological add-ons were reported in 4.9% of respondent farms. For those that did have them, the most common add-ons were in-parlour feeding systems (84.5%), automatic washers on the bulk tank (72.8%), ACRs (57.9%), entrance or exit gates controlled from the pit of the parlour (52.2%), and automatic washers on the milking machine (34.8%). Other add-ons included variable speed milk pumps (31.0%), a milk dump line (29.9%), variable speed vacuum drives (26.9%), electronic milk meters (23.6%), automatic drafting system (22.0%), automatic cluster flush (15.2%), automatic teat sprayers (14.9%), backing gates in the collecting yard (13.3%), automatic cow ID system (10.9%), dual vacuum or similar system for pulsation (alternatives to ACRs) (6.0%), non-electronic milk meters (5.7%), no technology add-ons (4.9%), automatic mastitis detection (4.1%), and automatic in-cluster dipping (1.9%).

Table 3 Average number of technology add-ons per parlour by herd size quartiles

Quartiles	Average number of technology add-ons
1 (< 73 cows)	3.0
2 (73–105 cows)	4.6
3 (106–159 cows)	5.7
4 (> 159 cows)	7.0
Average	5.1

(26.9%), electronic milk meters (23.6%), automatic drafting (22.0%), automatic cluster flush (ACF) (15.2%), automatic teat sprayers (14.9%), and backing gates in the collecting yard (13.3%). The least common parlour add-ons included automatic identification systems (10.9%), dual vacuum systems (6.0%), non-electronic milk meters (5.7%), automatic mastitis detection systems (4.1%), and automatic in-cluster dipping (1.9%).

Cluster disinfection occurred in 34.4% of herds. Of these, 59.1% conducted manual disinfection whilst 40.9% used an automated system of cluster disinfection. When manually done, 53.3% conducted it after every high cell count cow, 44.0% conducted it after every cow milked in the parlour and 2.7% did it “at the end of milking” or for “cows that recently had mastitis” or “during treatment of cows with mastitis”. The most commonly used cluster disinfection agents were peracetic acid-based products regardless of method of disinfection. Cluster disinfection products were in concentrate form that required addition




Fig. 5 Total % of parlour technological add-ons across all parlour types (n = 368)

to water and mixing before use on farm in 56.7% of respondent cases and ready-to-use products on 37.8% of respondent farms.

Milking management

Fore-milking was conducted in cases where clots were identified in the milk filter in the previous milking (56.6%), for cows that were freshly calved (46.1%), for cows showing signs of clinical mastitis (hot, swollen and painful udders) (41.6%), and when increases in bulk tank SCC occurred (36.5%). In addition, fore-milking was conducted by 13.9% of farmers at every milking, 8.0% reported they never do it and 7.5% reported to do it at every morning milking only. 'Other' responses included "after receiving milk recording results", "every morning during housing only", "once per week" and "[if] udder rub [is] abnormal" ('udder rub' referring to a practice whereby farmers gently massage the udder prior to milking to check for any palpable abnormalities and to stimulate milk let-down).

When asked where they strip the foremilk, 79.6% stripped it onto the floor, 14.9% into a gloved hand, 2.7% into a strip cup and 0.9% into a bare, non-gloved hand. CMT testing was utilised on 56.9% of respondent farms.

In terms of pre-milking preparation, 32.1% of respondents reported not doing any udder preparation. A dry wipe was carried out for cleaning the external surface of the teat prior to milking on 37.5% of respondent farms, a wash with a hose on 10.2%, individual udder wash cloths (defined as one cloth used per cow) on 2.7% and a communal wash cloth (defined as use of a single wash cloth on more than one cow) on 1.6% of farms. Less frequent methods of washing included the use of an udder brush (1.3%) and a pre-milking wash cup (1.1%). Pre-spraying was implemented on 18.9% of farms and pre-dipping on 7.0% of farms. Drying was conducted using an individual (one piece per cow) dry cloth or piece of paper towel on 10.8% of farms and using a communal dry cloth or piece of paper towel (one piece used on more than one cow) on 6.5% of respondent farms. Other options for pre-milking udder preparation included "dry wipe [in] winter only", "check for dirt and clean with hand", "none except soiled teats washed" and "pre-spray high SCC cows" or "pre-spray in springtime".

Post-milking teat disinfection most commonly involved spraying with a disinfectant solution, with 88.4% of farms using this method. Post-milking dipping occurred on 6.5% of farms and automatic in-cluster dipping on 0.5% of farms. No post-milking teat disinfection occurred on 3.0% of farms. 'Other' options (1.6%) included "post-spray in spring and autumn" and "spraying cows only when housed". The most commonly used teat disinfection products contained chlorhexidine or lactic acid as their

primary disinfectant agent; occasionally utilising a combination of the two with varying concentrations depending on brand.

Other udder hygiene practices involved clipping udders (18.8%), flaming udders (14.7%) and clipping tails (96.5%) in order to achieve better maintenance and ease of udder cleaning.

Gloves were always worn in the parlour on 81.6% of respondent farms. Gloves were sometimes worn in the parlour in 12.6% and never worn in 5.9% of responses. The most common type of gloves used were disposable gloves (85.2%), followed by reusable and washable gloves (8.1%).

SCC control

Mastitis treatment records were kept by 90.7% of respondents. Of these, 49.0% used a whiteboard, 48.7% used a farm book and 50.7% used a software application. 10.1% of respondents used another means of keeping mastitis treatment records. Given the nature of the checkbox question, participants could select more than one answer and therefore percentage figures equate to greater than 100%.

Collecting milk samples for bacteriological analysis was conducted rarely on 23.5% of respondent farms and never done on 15.5%. Reasons for doing it included the advice of a veterinarian (as opposed to being done autonomously) in 22.1% of cases, for both clinical and subclinical mastitis cases (22.4%), for clinical mastitis cases only (9.6%) and for subclinical mastitis cases only (6.9%). Advice on how to manage SCC was acquired from veterinarians by 76.9% of farmers, followed by discussion with colleagues (50.0%), processor milk quality advisors (39.2%) and websites (20.7%). Less frequently used sources of information on SCC management included other advisors (15.3%), magazines (14.5%), mastitis handbooks (11.6%) and on-site visits by mastitis experts (6.1%). Other sources of advice (2.4%) included "milk recording results" and "milking machine technician[s]". Again, given the nature of the checkbox question, participants could select more than one answer and therefore percentage figures equate to greater than 100%. The practicality of milk recording reports or text alerts for managing SCC was rated at a '4' or '5' of the five-point scale by 90.8% of the survey respondents, where '5' is a strong agreement to their usefulness.

When asked to choose what they considered most responsible for causing high SCC on farm, the most common answers were the milking parlour (33.2%), older cows in the herd (24.7%), housing conditions (16.6%) and the milking machine (9.9%). 'Milking parlour' refers to the general milking environment and associated milking management practices (e.g. parlour cleanliness, liner

condition, pre- and post- milking practices, etc.), whereas ‘milking machine’ refers to machine-specific functions which may influence source and spread of mastitis (e.g. vacuum and pulsation settings, cluster detachment timings, etc.). Less frequently chosen answers included freshly calved cows (1.6%) and grassland (0.3%). ‘Other’ answers (7.2%) included “general weakness for health”, “not milking the cows yourself”, “slow action by the farmer to a problem cow which lets the issue spread” and “selective dry cow therapy”. The source of high SCC was deemed unknown by 6.4% of respondents. In terms of managing high SCC cows, 78.3% of respondents milk them alongside the rest of the herd whilst 16.3% milk them after the rest of the herd.

Farmer demographics

Respondents were 96.8% male and 3.2% female. The most common age range was 45–54 (34.0%) followed by 35–44 (25.3%) and 55–64 (24.5%). Remaining groups included 10.9% aged between 25 and 34, 4.8% aged 65+ and 0.5% in the younger range of 18–24 years of age. Respondents were most commonly dairying for 20–30 years (26.6%), with 19.2% being in dairy for greater than 40 years and 5.6% for less than five years. Levels of education varied from primary (1.6%), to secondary (9.1%), to third level (23.0%) to various levels of courses completed. These included certificate courses in agriculture such as the ‘Green Cert’ (35.9%) and other courses in agricultural colleges that lasted for one year (12.9%) or greater than one year (12.6%).

For 2021, 40.7% respondents were very happy (score of ‘8’, ‘9’ or ‘10’ on the 10-point satisfaction scale) with their SCC management whilst 17.3% were very worried about it (score of ‘1’, ‘2’ or ‘3’ on the 10-point satisfaction scale). For 2022, 48.3% were very happy with their SCC management and 15.2% were worried. High levels of confidence that a low SCC (<200,000 cells/ml) is achievable on their farm was reported by 77.0% of respondents.

The effect of legislation changes on strict implementation of selective dry cow therapy was not predicted to affect the current drying-off practices of 34.2% of respondents (score of ‘8’, ‘9’ or ‘10’ on a 10-point scale), but was anticipated to greatly affect the practices of 22.5% of respondents (‘1’, ‘2’ or ‘3’ on a 10-point scale), based on their own responses. Confidence in the ability of selective dry cow therapy (SDCT) to control SCC was high (score of ‘8’, ‘9’ or ‘10’ on a 10-point scale) for 38.7% of respondents and low (‘1’, ‘2’ or ‘3’ on a 10-point scale) for 28.1% of respondents. For farmers already implementing SDCT, dry-off treatment with teat sealant alone was implemented based on individual cow SCC records on 62.9% of farms, on available records of clinical cases and their outcomes throughout the lactation for 54.5% of

farms, on individual cow factors for 28.4%, on milk yield records for 22.2%, and on the results of individual CMT testing on 16.5% of farms. None of the above were used in decision making for 8.4% of farms. The most common SCC cut-off point for using teat sealant only dry-off treatments was 100,000 cells/ml (86.1%).

Satisfaction with their career in dairy was scored highly (‘8’, ‘9’ or ‘10’ on a 10-point scale) by 73.4% of respondents and less favourably (‘1’, ‘2’ or ‘3’) by 5.4% of survey respondents. The most common answer was a score of ‘10’ with 29.4% of respondents claiming immense fulfilment from their profession.

Discussion

A study by Palma Molina et al. [23] documented the prevalence of certain technologies but did not examine milk quality on farms. Other Irish studies document certain aspects of management practices and their relationship with BTSCC [24, 35, 36], but did not tie these together with parlour technologies. This study documented the farm and parlour management practices, parlour technologies, milking management practices, SCC control strategies and farmer demographics on a sample of commercial Irish dairy farms, thereby filling current gaps in the literature. The geographical distribution of responses in this study aligns with the distribution of milk producers in Ireland. Most responses were in Munster (59.0%), particularly in Cork, the county with the greatest number of dairy cows according to the Central Statistics Office (CSO) [11]. According to the 2022 Teagasc National Farm survey [37], 72% of Irish dairy animals were in the south of the country. The national average herd size in 2022 was 93 cows [37], meaning our study has an inherent inference bias towards larger herds (average herd size of 131 cows). Herd size quartiles in this study reflect a smaller average herd size per quartile than other studies, such as those reported by Hogan et al. [38]. Cyclical, seasonal trends of BTSCC found in this study are similar to those reported by O’Connell et al. [34] for both calving patterns and herd size quartiles.

Parlour technologies and management

Good facilities and modernization of dairy operations have been shown in the literature to correlate with improved production, efficiency and personal satisfaction [39]. In line with this, dairy farmers worldwide are becoming increasingly interested in the implementation of technology to replace labour and to improve efficiency [40–42]. Our study found that 46.8% of dairy parlours were less than 10 years of age, with 20.1% being less than five years of age. This could reflect growing dairy cow numbers on these particular farms which therefore necessitated increased space to accommodate them, or

could be indicative of a need for modernization to suit current milking efficiency or labour demands. Despite this finding, 63.2% of respondents reported not implementing any major upgrades in the last five years. Though we can assume the new parlours fall under the 36.8% who did, we cannot verify this. Our study found that smaller herds had lower levels of technology add-ons (except for dual vacuum systems and automatic in-cluster dipping; Supplementary Table 4). The use of automation increased with herd size quartile (Table 3), for example herds in Q4 had an average of 7.0 technology add-ons compared to an average of 3.0 for Q1 herds (Table 3). However, similar levels of uptake of dual vacuum systems, automatic teat sprayers and non-electronic milk meters was noted across all four quartiles, as was increased implementation of automatic in-cluster dipping in lower quartile (Q1 and Q2) herds compared to no uptake in Q3 and Q4 respondent herds. The most common technological additions on respondent farms were in-parlour feeding systems (84.5%), automatic washers on the bulk tank (72.8%), ACRs (57.9%), and entrance or exit gates controlled from the pit of the parlour (52.2%). In this study, AMS parlours and rotaries had more parlour technologies on average than other types of parlour, with an average of 12.3 and 10.8 additions respectively (Supplementary Table 5). This finding was in agreement with the work of Yang et al. [43] which found that New Zealand rotary parlours were associated with a significantly increased number of automated labour-saving and data-capture technologies compared to herringbones. A recent study by Prendergast et al. [44] demonstrated that the number of automations on herringbone farms had a strong positive correlation with milking efficiency in terms of cow throughput and litres of milk harvested per hour, but a negligible correlation was documented between milking efficiency and automations for rotary parlours; likely due to minimal variation in automation presence on these farms. Our study also supports the Yang et al. [43] findings that ACRs and in-parlour feeding were among the most prevalent technological additions in both herringbones (34.9–66.3% and 53–78.3% respectively depending on the type of herringbone) and rotaries (100% for both respectively), though those in the Yang et al. [43] study had a much higher rate of automatic teat sprayer use than we found.

Another parlour management finding in this study was that 34.4% of survey respondents reported changing their liners only once per year. The current advice for spring-calving herds is to change liners every 2,000 milkings or every 6 months, whichever comes first [10]. This is true particularly for commonly encountered nitrile rubber liners, though silicone liners have been described in literature as having greater longevity

of 5000 milkings or more [45, 46]. The liner is the only element of the milking machine that is in direct contact with the cow's teat, therefore it is important that they are operating effectively. When liners are worn, they lose their shape and do not massage the teat correctly, resulting in longer milking times and reduced yields [47]. Worn liners are also a source of bacteria, particularly contagious mastitis-causing strains, as the cracks in the liner surface provide ideal environments for bacterial accumulation and growth [48].

Milking management

Hygiene and stringent management practices during milking are linked with reduced farm-level SCC [24, 26, 34, 49, 50]. Pre-milking routines generally consist of stimulating milk let-down and preparing teats for cluster attachment. Regulation (EC) No 853/2004 states that raw milk should only be collected from cows which “present no sign of disease that might result in the contamination of milk... or a recognisable inflammation of the udder” [51]. Our study found that 13.9% of respondent farms conducted fore-milking as part of their daily milking routine and 8.0% reportedly never conduct it. This low prevalence of consistent fore-milking may inhibit farmers' ability to identify any visible abnormalities in the milk such as blood or clots, which indicate evidence of clinical mastitis (CM) [52], especially given the low incidence of automatic mastitis detection systems (4.1%). Similarly, fore-milking is considered an effective stimulus for milk let-down [53]. Other pre-milking management practices include washing and drying of teats prior to cluster attachment. Clusters must be attached to clean, dry teats in order to minimise risk of environmental mastitis. However, we found that 32.2% of respondent farmers do not do any form of pre-milking teat preparation. Washing udders prior to milking, conducted by 14.5% of respondent farmers using either a hose or a wash cloth, is generally discouraged in the absence of adequate drying afterwards as pathogens can be distributed all over the surface of the udder and teats [30]. Should washing be required, only the teats should be washed, using minimal water, and should be thoroughly dried afterwards [52]. Galton et al. [54] demonstrated that washing and subsequent drying of teats resulted in low bacterial counts on teat ends and therefore less risk of new mastitis infection. Pankey et al. [55] showed that inadequate cleaning and drying of teats caused environmental mastitis pathogen numbers in milk to increase. Similarly, drying of teats via communal means, such as was reported by 6.5% of survey respondents, is strongly discouraged due to risk of mastitis pathogen spread from cow to cow and was associated with a higher monthly rate of clinical mastitis in a study by Ruegg [56]. Post-milking teat disinfection is required

to destroy pathogens on teat skin at the end of milking. Only 3.0% of respondent farms reported no post-milking teat disinfection which, though worrying, were considerably outweighed by those conducting spraying (88.4%) and dipping (6.5%) of teats. Spraying offers an advantage in terms of time saving and more limited handling of cows that are particularly restless in the parlour or are first-time milkers [57]. Appropriate coverage is essential [58], with AHI Cell Check guidelines [59] suggesting a minimum of 10 ml of dip or 15 ml of spray per cow, per milking.

SCC control

SCC and mastitis management practices were also investigated in this study. Almost six percent (5.9%) of farmers reported never wearing gloves in the parlour for milking, despite evidence in literature that glove-wearing practices are associated with reduced risk of spreading mastitis-causing bacteria from cow to cow via milkers' hands [60, 61]. Persistently high SCC and recurrent incidence of CM were two of the greatest indicators for culling in this study. Lack of consistent fore-milking and also lack of engagement with bacteriology of milk samples, conducted 'rarely' or 'never' on 39.0% of respondent farms, could contribute to increased BTSCC through slower identification of clinical cases. A well-managed combination of hygiene practices and suitable antibiotic administration has long been proven to provide superior mastitis control to implementation of either measure alone [29]. Though the majority (77.0%) of respondents in our study strongly agreed that they believe a low SCC is attainable on their farms, 51.7% reported that amendments to the legislation surrounding SDCT will change their current management practices around drying-off and 28.1% reported concurring low levels of confidence in their ability to maintain a low SCC in the advent of said changes. These legislative changes refer to the new Veterinary Medicines Regulation EU 2019/6 [62] which came into effect in January 2022. This legislation specifically prohibits prophylactic antibiotic use, other than in well-defined cases for an individual or a restricted number of animals when the risk of infection is very high and the consequences are likely to be severe (Article 107 [3]). National and farm-level recommendations "in support of improved mastitis control and intramammary antimicrobial stewardship in the Irish dairy industry" have been created by AHI Cell-Check [28]. Farmers in our study found that support offered by milk recording reports and texts were useful, as were interactions with their veterinary professionals and milk quality advisors. This is an important finding, as fostering strong working relationships between

farmers and their veterinary professionals, in particular, is more important than ever in order to ensure good antimicrobial stewardship and maintenance of optimal herd health. A novel approach to designing behaviour change interventions for antimicrobial use in Irish agriculture is described by Regan et al. [63]. Further factors which can influence behaviour in relation to antimicrobial use are outlined in an international study by McKernan et al. [64].

Farmer demographics

General attitudes assessed in this study showed that farmers were generally quite content with their career in dairy. We found that 29.4% of respondents claimed immense fulfilment from their profession; awarding their love for their career a score of '10' out of '10' on a satisfaction scale. However, other studies have shown that there are considerable concerns in the dairy sector surrounding recruitment and retention of labour [65]. The intense seasonal workload associated with pasture-based farming [18, 38] can lead to increased levels of stress [66], mental health problems [67] and difficulties in sustaining an adequate quality of life [68]. Our study found that 63.3% of survey respondents were over the age of 45 and 66.1% of respondents had sustained a career in dairy for over 20 years, with 19.2% of these having farmed for over 40 years. Greater emphasis on incentives such as a good work-life balance and increased family time [69], and creation of a more sustainable workload are essential in enticing the younger generation towards a career in farming [70]. This supports recent research carried out by Huey et al. [71] and Regan et al. [72]. Moving forward, further efforts in offering farmers support and decision-making tools for farm and parlour management, as well as continued education for the professionals they most engage with to ensure there is consistency in advice and methods of advice dissemination, could be of great benefit to the Irish dairy industry.

Study limitations

We are aware of a level of bias pertaining to the fact that respondent herd sizes are larger than the national average. It is important to note that this study was based on self-reported management practices, behaviour and attitudes of farmers. The farmers participating in this study could be different than the average Irish farmer because of their willingness to self-select participation in the survey (inferring a certain level of selection bias). There is also a chance that farmers reported socially desirable answers which could directly add bias to the results. It is important to note that, although the survey was extensive and developed alongside experts in the field of mastitis, the total dataset of parlour technologies, management

practices, attitudes and behaviour regarding milking management may be incomplete.

Conclusion

In this study, we successfully filled current knowledge gaps around farm management practices, parlour management including rates of technology adoption, general milking management, SCC control strategies and farmer attitudes and behaviour towards certain SCC management practices by completing a survey of Irish dairy farms. Our study found that the most common parlour technology additions were in-parlour feeders (84.5%), automatic washers on the bulk tank (72.8%) and ACRs (57.9%). Rotary parlours and AMS parlours had the most parlour technologies of any parlour type (10.8 and 12.3 add-ons on average respectively). Technology add-ons were less prevalent in smaller herds (average number for Q1 being 3.0 add-ons per herd compared to 7.0 for Q4 herds), except for dual vacuum systems and automatic in-cluster dipping. Despite finding areas for improvement with regard to frequency of liner changes (51.5% changing at the recommended intervals of every 2000 milkings or every six months), glove-wearing practices (18.5% wearing them sometimes or never), using a communal means of drying teats in the parlour (6.5% using a communal cloth or paper towel to dry teats prior to milking), and engagement with bacteriology of milk samples (rarely or never done on 39.0% of farms), we also found evidence of high levels of good management practices, including the documentation of mastitis treatment records (90.7%) and high use of post-milking teat disinfection (94.9% either dipping or spraying). We discovered that Irish dairy farmers are relatively content in their careers but face pressures following changes to the legislation regarding antimicrobial use in their herds. However, there are high levels of confidence that low herd level SCC is achievable on respondent farms. Farmers in our study found that support offered by milk recording reports and texts were useful for reducing their BTSCC, as were interactions with their veterinary professionals and milk quality advisors.

Abbreviations

SCC	Somatic cell count
AMS	Automatic milking system
BTSCC	Bulk tank somatic cell count
ACR	Automatic cluster remover
PLF	Precision livestock technology
CMT	California Mastitis Test
AHI	Animal Health Ireland
ICBF	Irish Cattle Breeding Federation
ACF	Automatic cluster flush
SDCT	Selective dry cow therapy
CSO	Central Statistics Office
CM	Clinical mastitis

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13620-024-00267-y>.

Supplementary Material 1.

Supplementary Material 2.

Acknowledgements

The authors would like to acknowledge the help and support offered by the AHI CellCheck Technical Working Group for their contributions to the development of the survey. We would also like to acknowledge the support of ICBF and their facilitating data transfers to this study. We extend gratitude to each of the milk processors for distributing this survey and to each of the respondent farmers for taking the time to complete the survey and contribute to this research. We would also like to thank Joe Patton of the Teagasc advisory services for his support.

Authors' contributions

AB assisted in the structuring of the survey and through guidance prior to the cognitive interviews. AUIC, JU, PSB, ER, CMC and CMC were involved in the development of survey questions. AUIC, JU and PSB analysed the survey data. AUIC led the writing of the manuscript. All authors contributed to the writing and editing of the final manuscript.

Funding

This publication has emanated from research conducted with the financial support of Science Foundation Ireland (SFI) and the Department of Agriculture, Food and Marine on behalf of the Government of Ireland through the VistaMilk centre under Grant Number [16/RC/3835].

Availability of data and materials

The participants of this study did not give written consent for their data to be shared publicly, so due to the sensitive nature of the research supporting data is not available.

Declarations

Ethics approval and consent to participate

This study was granted an ethics exemption from the University College Dublin (UCD, Belfield, Dublin 4) AREC ethics committee (reference number AREC-E-22-11-Ryan).

Mandatory consent was acquired for this from all respondents at the beginning of the survey. Without granting consent, farmers were unable to access the survey.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Teagasc, Animal & Grassland Research and Innovation Centre, Moorepark, Fermoy, Co. Cork P61 C997, Ireland. ²School of Veterinary Medicine, University College Dublin, Belfield D04 W6F6, Dublin 4, Ireland. ³Animal Health Ireland, 2-5 The Archways, Carrick On Shannon N41 WN27, Co. Leitrim, Ireland.

Received: 22 November 2023 Accepted: 6 March 2024

Published online: 06 May 2024

References

1. (Ibec) DII. Irish Dairy Industry Economic & Social Snapshot. [cited 2023 Aug 15]. <https://www.ibec.ie/dairyindustryireland/our-dairy-story/economics-and-social#:~:text=The dairy industry is a,employment focused in rural areas>

2. McCoy F. Mastitis Control Programmes - The Irish Experience. In: Proceedings of the British Mastitis Conference, Sixways, Worcester. 2012:65–9.
3. Halasa T, Huijps K, Østerås O, Hogeveen H. Economic effects of bovine mastitis and mastitis management: A review. *Vet Q*. 2007;29(1):18–31. <https://doi.org/10.1080/01652176.2007.9695224>.
4. Seegers H, Fourichon C, Beaudeau F. Production effects related to mastitis and mastitis economics in dairy cattle herds. *Vet Res*. 2003;34(5):475–91. <https://doi.org/10.1051/vetres:2003027>
5. European Union. European Council Directive 92/46/EEC of 16 June 1992. Laying down the health rules for the production and placing on the market of raw milk, heat-treated milk and milk-based products. *Off J Eur Communities*. 1992;35(L268):1–32.
6. DAFM. EU hygiene regulations in relation to TBC and SCC for raw milk. Department of agriculture, food and the marine. Trader notice DH/TN/01/2018 (revised May 2018) 2018 p. 1–4. <https://www.gov.ie/pdf/?file=https://assets.gov.ie/246237/6152ce68-e82b-4106-acd8-a827f00b0055.pdf>.
7. DeLong KL, Lambert DM, Schexnayder S, Krawczel P, Fly M, Garkovich L, et al. Farm business and operator variables associated with bulk tank somatic cell count from dairy herds in the southeastern United States. *J Dairy Sci*. 2017;100(11):9298–310. <https://doi.org/10.3168/jds.2017-12767>.
8. Deluyker HA, Gay JM, Weaver LD. Interrelationships of Somatic Cell Count, Mastitis, and Milk Yield in a Low Somatic Cell Count Herd. *J Dairy Sci*. 1993;76(11):3445–52.
9. More SJ, Madden JM, McAloon CI. Prescribing and sales of intramammary antimicrobials in Ireland in 2019 and 2020: the role of milk purchasers. *Ir Vet J*. 2022;75(1):20. <https://doi.org/10.1186/s13620-022-00227-4>.
10. O'Brien B. Moorepark Dairy Levy Research Update. 2008 [cited 2023 Jul 25]. Milk Quality Handbook: Practical Steps to Improve Milk Quality. <http://hdl.handle.net/11019/1164>
11. Central Statistics Office (CSO). Livestock Survey December 2022. 2022 [cited 2023 Jul 25]. [https://www.cso.ie/en/releasesandpublications/ep/p-lsd/livestocksurveydecember2022/#:~:text=Statistician's Comment&text=%22While the total number of,1.51 million \(%2B0.3%25\).](https://www.cso.ie/en/releasesandpublications/ep/p-lsd/livestocksurveydecember2022/#:~:text=Statistician's Comment&text=%22While the total number of,1.51 million (%2B0.3%25).)
12. Central Statistic Office (CSO). Milk Statistics December 2022. 2022 [cited 2023 Jul 25]. <https://www.cso.ie/en/releasesandpublications/er/ms/milkstatisticsdecember2020/%0Ainternal-pdf/651/milkstatisticsdecember2020.html>
13. Kelly P, Dillon P. Irish dairying – rapid expansion, structural change and future plans. In: Moorepark dairy levy research update, international agricultural workforce conference, Animal and Grassland Research and Innovation Centre (AGRICE). 2018. p. 25–7. https://www.teagasc.ie/media/website/publications/2018/People_in_dairying_-_Kelly.pdf.
14. O'Brien B. Mastitis and SCC. 2016 [cited 2023 Jul 25]. p. 183–90. <https://www.teagasc.ie/media/website/animals/dairy/MilkQandMastitis.pdf>
15. O'Brien B, Shalloo L, O'Donnell S, Butler AM, Gleeson D, O'Donovan K. Labour Productivity – Effects of Scale, Capital Investment and Adoption of Novel Technology. In: Proceedings from the 16th International Farm Management Association Congress. 2007. p. 407–507. https://www.researchgate.net/profile/David-Gleeson/publication/267960290_labour_productivity_effects_of_scale_capital_investment_and_adoption_of_novel_technology/links/54b648680cf2318f09a2d7c/labour-productivity-effects-of-scale-capital-investment-and-
16. Timlin M, Tobin JT, Brodtkorb A, Murphy EG, Dillon P, Hennessy D, et al. The impact of seasonality in pasture-based production systems on milk composition and functionality. *Foods*. 2021;10(3):607. <https://doi.org/10.3390/foods10030607>.
17. O'Donovan K, O'Brien B, Ruane D, Kinsella J, Gleeson D. Labour Input on Irish Dairy Farms and the Effect of Scale and Seasonality. *J Farm Manag*. 2008;13(5):38–53. https://www.iagrm.com/content/large/journals/jofm/volume_13/jofm-vol13-no5-pages-346-361.pdf
18. Deming J, Gleeson D, O'Dwyer T, Kinsella J, O'Brien B. Measuring labor input on pasture-based dairy farms using a smartphone. *J Dairy Sci*. 2018;101(10):9527–43. <https://doi.org/10.3168/jds.2017-14288>.
19. Shortall J, Shalloo L, Foley C, Sleator RD, O'Brien B. Investment appraisal of automatic milking and conventional milking technologies in a pasture-based dairy system. *J Dairy Sci*. 2016;99(9):7700–13. <https://doi.org/10.3168/jds.2016-11256>.
20. Hogan C, Kinsella J, O'Brien B, Markey A, Beecher M. Estimating the effect of different work practices and technologies on labor efficiency within pasture-based dairy systems. *J Dairy Sci*. 2022;105(6):5109–23. <https://doi.org/10.3168/jds.2021-21216>.
21. Galton DM. Effects of an automatic postmilking teat dipping system on new intramammary infections and iodine in milk. *J Dairy Sci*. 2004;87(1):225–31. [https://doi.org/10.3168/jds.S0022-0302\(04\)73161-6](https://doi.org/10.3168/jds.S0022-0302(04)73161-6).
22. Ohnstad I, Olde Riekerink RGM, Hogewerf P, de Koning CAJM, Barkema HW. Short communication: Effect of automatic postmilking teat disinfection and cluster flushing on the milking work routine. *J Dairy Sci*. 2012;95(5):2567–70. <https://doi.org/10.3168/jds.2011-4946>
23. Palma-Molina P, Hennessy T, O'Connor AH, Onakuse S, O'Leary N, Moran B, et al. Factors associated with intensity of technology adoption and with the adoption of 4 clusters of precision livestock farming technologies in Irish pasture-based dairy systems. *J Dairy Sci*. 2023;106(4):2498–509. <https://doi.org/10.3168/jds.2021-21503>.
24. Kelly P, O'Sullivan K, Berry D, More S, Meaney W, O'Callaghan E, et al. Farm management factors associated with bulk tank somatic cell count in Irish dairy herds. *Ir Vet J*. 2009;62(S4):S45. <https://doi.org/10.1186/2046-0481-62-S4-S45>.
25. Barkema HW, Van Der Ploeg JD, Schukken YH, Lam TJGM, Benedictus G, Brand A. Management style and its association with bulk milk somatic cell count and incidence rate of clinical mastitis. *J Dairy Sci*. 1999;82(8):1655–63. [https://doi.org/10.3168/jds.S0022-0302\(99\)75394-4](https://doi.org/10.3168/jds.S0022-0302(99)75394-4).
26. Hutton CT, Fox LK, Hancock DD. Mastitis Control Practices: Differences Between Herds with High and Low Milk Somatic Cell Counts. *J Dairy Sci*. 1990;73(4):1135–43. [cited 2022 Mar 16]. [https://doi.org/10.3168/jds.S0022-0302\(90\)78774-7](https://doi.org/10.3168/jds.S0022-0302(90)78774-7)
27. Jones T, Ohnstad I. Milking procedures recommended for the control of bovine mastitis. In *Pract*. 2002;24(9):502–11. <https://doi.org/10.1136/inpract.24.9.502>.
28. More SJ, McAloon C, Silva Boloña P, O'Grady L, O'Sullivan F, McGrath M, et al. Mastitis Control and Intramammary Antimicrobial Stewardship in Ireland: Challenges and Opportunities. *Front Vet Sci*. 2022;9(April). <https://doi.org/10.3389/fvets.2022.748353>
29. Philpot WN. Control of Mastitis by Hygiene and Therapy. *J Dairy Sci*. 1979;62(1):168–76. [https://doi.org/10.3168/jds.S0022-0302\(79\)83218-X](https://doi.org/10.3168/jds.S0022-0302(79)83218-X).
30. Neave FK, Dodd FH, Kingwill RG, Westgarth DR. Control of Mastitis in the Dairy Herd by Hygiene and Management. *J Dairy Sci*. 1969/05/01. 1969;52(5):696–707. [https://doi.org/10.3168/jds.S0022-0302\(69\)86632-4](https://doi.org/10.3168/jds.S0022-0302(69)86632-4)
31. Jansen J, van den Borne BHP, Renes RJ, van Schaik G, Lam TJGM, Leeuwis C. Explaining mastitis incidence in Dutch dairy farming: The influence of farmers' attitudes and behaviour. *Prev Vet Med*. 2009;92(3):210–23. <https://doi.org/10.1016/j.prevetmed.2009.08.015>.
32. Dillman D, Smyth J, Christian L. Internet, Phone, Mail, and Mixed-Mode Surveys: the Tailored Design Method. Fourth ed. (Fourth edition) Wiley. Hoboken, New Jersey: Wiley; 2014.
33. Central Statistic Office (CSO). Census of Agriculture 2020 Detailed Results. [cited 2023 Jul 25]. <https://www.cso.ie/en/releasesandpublications/ep/p-coa/censusofagriculture2020detailedresults/standardoutput/>
34. O'Connell A, McParland S, Ruegg PL, O'Brien B, Gleeson D. Seasonal trends in milk quality in Ireland between 2007 and 2011. *J Dairy Sci*. 2015;98(6):3778–90. <https://doi.org/10.3168/jds.2014-9001>.
35. McAloon CI, McCoy F, More SJ. Trends in estimated intramammary antimicrobial usage in the Irish dairy industry from 2003 to 2019. *JDS Commun*. 2021;2(5):271–6. <https://doi.org/10.3168/jdsc.2021-0081>.
36. Clabby C, McParland S, Dillon P, Arkins S, Flynn J, Murphy J, et al. Internal teat sealants alone or in combination with antibiotics at dry-off – the effect on udder health in dairy cows in five commercial herds. *Animal*. 2022;16(2): 100449. <https://doi.org/10.1016/j.animal.2021.100449>.
37. Dillon E, Donnellan T, Moran B, John L. Teagasc National Farm Survey 2022. 2022 [cited 2023 Jul 25]. https://www.teagasc.ie/media/website/publications/2023/NFS_prelim_results_2022.pdf
38. Hogan C, Kinsella J, O'Brien B, Gorman M, Beecher M. An examination of labor time-use on spring-calving dairy farms in Ireland. *J Dairy Sci*. 2022;105(7):5836–48. <https://doi.org/10.3168/jds.2022-21935>.
39. Bewley J, Palmer RW, Jackson-Smith DB. An overview of experiences of wisconsin dairy farmers who modernized their operations. *J Dairy Sci*. 2001;84(3):717–29. [https://doi.org/10.3168/jds.S0022-0302\(01\)74526-2](https://doi.org/10.3168/jds.S0022-0302(01)74526-2).
40. Wilson P. Decomposing variation in dairy profitability: the impact of output, inputs, prices, labour and management. *J Agric Sci*. 2011;149(4):507–17. <https://doi.org/10.1017/S0021859610001176>.
41. Dela Rue BT, Eastwood CR, Edwards JP, Cuthbert S. New Zealand dairy farmers preference investments in automation technology over decision-support technology. *Anim Prod Sci*. 2020;60(1):133. <https://doi.org/10.1071/AN18566>.

42. Edwards JP, Dela Rue BT, Jago JG. Evaluating rates of technology adoption and milking practices on New Zealand dairy farms. *Anim Prod Sci*. 2015;55(6):702. <https://doi.org/10.1071/AN14065>.
43. Yang W, Edwards JP, Eastwood CR, Dela Rue BT, Renwick A. Analysis of adoption trends of in-parlor technologies over a 10-year period for labor saving and data capture on pasture-based dairy farms. *J Dairy Sci*. 2021;104(1):431–42. <https://doi.org/10.3168/jds.2020-18726>.
44. Prendergast R, Murphy MD. The effects of seasonality, management, infrastructure and automation on the milking efficiency of herringbone and rotary milking parlors in Ireland. *J Dairy Sci*. 2023. <https://doi.org/10.3168/jds.2023-23540>.
45. Mein G, Reinemann D, O'callaghan E, Ohnstad I. Where the rubber meets the teat and what happens to milking characteristics. *Proc 100 years with Liners Pulsators*. 2003;(October 2003):431–8.
46. de Koning K, Rønningen O, Bjerring M, Ipema A. Testing of liners and measuring material properties. In: *IDF World Dairy Summit & Centenary '100 years with Liners and Pulsators in Machine Milking'*. 2003.
47. Davis MA, Reinemann DJ. Effect of liner design on milking characteristics. In: *Proceedings, 39th AABP-NMC International Symposium on Mastitis and Milk Quality*, Vancouver, BC, Canada. 2000. https://www.researchgate.net/publication/237576275_effect_of_liner_age_on_milking_characteristics#fullTextFileContent
48. Fox LK, Gershman M, Hancock DD, Hutton CT. Fomites and reservoirs of *Staphylococcus aureus* causing intramammary infections as determined by phage typing: the effect of milking time hygiene practices. *Cornell Vet*. 1991;81(2):183–93. <http://www.ncbi.nlm.nih.gov/pubmed/2029841>
49. Barkema HW, Schukken YH, Lam TJGM, Beiboer ML, Benedictus G, Brand A. Management practices associated with low, medium, and high somatic cell counts in bulk milk. *J Dairy Sci*. 1998;81(7):1917–27. [https://doi.org/10.3168/jds.S0022-0302\(98\)75764-9](https://doi.org/10.3168/jds.S0022-0302(98)75764-9).
50. Peeler EJ, Green MJ, Fitzpatrick JL, Morgan KL, Green LE. Risk factors associated with clinical mastitis in low somatic cell count British dairy herds. *J Dairy Sci*. 2000;83(11):2464–72. [https://doi.org/10.3168/jds.S0022-0302\(00\)75138-1](https://doi.org/10.3168/jds.S0022-0302(00)75138-1).
51. European Commission. Regulation (EC) N° 853/2004 of the European Parliament and of the Council of 29 April 2004 laying down specific hygiene rules for on the hygiene of foodstuffs. *Off J Eur Union*. 2004;L 139(853):55.
52. Rasmussen MD. A Review of Milking Preparation: the Science. In: *National Mastitis Council Ann Meet*. 2000. p. 104–10.
53. Bruckmaier RM, Blum JW. Oxytocin Release and Milk Removal in Ruminants. *J Dairy Sci*. 1998;81(4):939–49. [https://doi.org/10.3168/jds.S0022-0302\(98\)75654-1](https://doi.org/10.3168/jds.S0022-0302(98)75654-1).
54. Galton DM, Petersson LG, Merrill WG, Bandler DK, Shuster DE. Effects of Premilking Udder Preparation on Bacterial Population, Sediment, and Iodine Residue in Milk. *J Dairy Sci*. 1984;67(11):2580–9. <https://linkinghub.elsevier.com/retrieve/pii/S0022030284816161>
55. Pankey JW, Wildman EE, Drechsler PA, Hogan JS. Field trial evaluation of premilking teat disinfection. *J Dairy Sci*. 1987;70(4):867–72. [https://doi.org/10.3168/jds.S0022-0302\(87\)80085-1](https://doi.org/10.3168/jds.S0022-0302(87)80085-1).
56. Ruegg PL. Pre-milking cow preparation - secret methods of producing high quality milk. In: *NMC Regional Meeting Proceedings (2004)*. 2004. p. 34–40.
57. Meaney WJ. Post-Milking Teat Disinfection: A Comparison of Dipping and Spraying. . Vol. 13, *Irish Journal of Agricultural Research*. 1974. <https://www.jstor.org/stable/25555721>
58. Pankey JW, Watts JL. Evaluation of spray application of postmilking teat sanitizer. *J Dairy Sci*. 1983;66(2):355–8. [https://doi.org/10.3168/jds.S0022-0302\(83\)81797-4](https://doi.org/10.3168/jds.S0022-0302(83)81797-4).
59. Animal Health Ireland. Cell Check Farm Guidelines For Mastitis Control : Lactation, Guideline 07. 2021; https://animalhealthireland.ie/assets/uploads/2021/06/FarmGuidlines_-_V6_001_WebVersion_Lactation_Guideline_07.pdf
60. Olde Riekerink RGM, Sampimon OC, Eerland VJ, Swarts MJ, Lam TJGM. Mastitis control: From science to practice. . Lam TJGM, editor. *Mastitis Control: From Science to Practice*. The Netherlands: Wageningen Academic Publishers; 2008. 77–82 p. <https://doi.org/10.3920/978-90-8686-649-6>
61. Jansen J, Renes RJ, Lam TJGM. Mastitis Control: take up the gloves! In: *International Symposia on Veterinary Epidemiology and Economics proceedings, ISVEE 12: Proceedings of the 12th Symposium, Durban, South Africa, Theme 1 - Surveillance and disease control: Mastitis, Disease monitoring & control, Evaluation of animal disease*. Durban, South Africa; 2009. p. 472–4. https://www.researchgate.net/publication/41042356_Mastitis_Control_take_up_the_gloves.
62. Regulation (EU) 2019/6. Regulation (EU) 2019/6 of the European Parliament and of the Council of 11 December 2018 on the Manufacture, Placing on the Market and Use of Medicated Feed, Amending Regulation (EC) No 1831/2003 of the European Parliament and of the Council and Repealing. *Off J Eur Union*. 2019;L4(726):43–167.
63. Regan Á, Burrell A, McKernan C, Martin H, Benson T, McAloon C, et al. Behaviour change interventions for responsible antimicrobial use on farms. *Ir Vet J*. 2023;76(1):1–13. <https://doi.org/10.1186/s13620-023-00236-x>.
64. McKernan C, Benson T, Farrell S, Dean M. Antimicrobial use in agriculture: critical review of the factors influencing behaviour. *JAC-Antimicrobial Resist*. 2021;3(4). <https://doi.org/10.1093/jacamr/dlab178>
65. Eastwood CR, Greer J, Schmidt D, Muir J, Sargeant K. Identifying current challenges and research priorities to guide the design of more attractive dairy-farm workplaces in New Zealand. *Anim Prod Sci*. 2020;60(1):84. <https://doi.org/10.1071/AN18568>.
66. Brennan M, Hennessy T, Meredith D, Dillon E. Weather, Workload and Money: Determining and Evaluating Sources of Stress for Farmers in Ireland. *J Agromedicine*. 2022;27(2):132–42. <https://doi.org/10.1080/1059924X.2021.1988020>.
67. Lunner Kolstrup C, Kallioniemi M, Lundqvist P, Kymäläinen HR, Stallones L, Brumby S. International Perspectives on Psychosocial Working Conditions, Mental Health, and Stress of Dairy Farm Operators. *J Agromedicine*. 2013;18(3):244–55. <https://doi.org/10.1080/1059924X.2013.796903>.
68. Contzen S, Häberli I. Exploring dairy farmers' quality of life perceptions – A Swiss case study. *J Rural Stud*. 2021;88:227–38. <https://doi.org/10.1016/j.jrurstud.2021.11.007>.
69. Russell RA, Bewley JM. Characterization of Kentucky dairy producer decision-making behavior. *J Dairy Sci*. 2013;96(7):4751–8. <https://doi.org/10.3168/jds.2012-6538>.
70. Beecher M, Gorman M, Kelly P, Horan B. Careers in dairy: adolescents perceptions and attitudes. *J Agric Educ Ext*. 2019;25(5):415–30. <https://doi.org/10.1080/1389224X.2019.1643745>.
71. Huey S, Kavanagh M, Regan A, Dean M, McKernan C, McCoy F, et al. Engaging with selective dry cow therapy: understanding the barriers and facilitators perceived by Irish farmers. *Ir Vet J*. 2021;74(1):28. <https://doi.org/10.1186/s13620-021-00207-0>
72. Regan Á, Burrell A, McKernan C, Martin H, Benson T, McAloon C. Use of antimicrobials in animal health on the island of Ireland: knowledge, attitudes and behaviour. *SafeFood AMU*. 2022; <https://www.safeFood.net/getattachment/def7240c-1abc-4092-a79a-bd9841be1b6d/AMR.pdf?lang=en-IE>

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.